

Insecticide Treatment Options for Western Bean Cutworm in Field Corn

Updated December 06, 2019

See <https://www.greenbook.net> or other resources for additional registrations.

*Consult product label for specific information and restrictions.

IRAC Mode of Action Classification:

- Group 1 = Acetylcholine esterase inhibitors; 1A = Carbamates, 1B = Organophosphates
- Group 3 = Sodium channel modulators
- Group 5 = Nicotinic acetylcholine receptor agonists (allosteric), Spinosyns
- Group 18 = Ecdysone agonists/moulting disruptors; 18A = Diacylhydrazines
- Group 22 = Voltage-dependent sodium channel blockers, 22A = Oxadiazines
- Group 28 = Ryanodine receptor modulators, Diamides

R = Restricted-Use Product

Mode Of Action	Product Name	Common Name	Rate (Formulation/Acre)	Restrictions/Comments
3 R	Ambush 2EC	permethrin	3.2-6.4 fl oz	PHI 30 days for grain or fodder (stover). Forage may be harvested on the day of application.
3 R	Arctic 3.2EC	permethrin	2-4 fl oz	PHI 30 days for harvest of grain or fodder (stover). Forage may be harvested on the day of application.
3 R	Asana XL	esfenvalerate	2.9-5.8 fl oz	PHI 21 days. Apply before larvae enter the ear.
3 R	Baythroid XL, Tombstone	cyfluthrin	1.6-2.8 fl oz	PHI 21 days for grain or fodder and 0 days for green forage.
28	Belt SC	flubendiamide	2-3 fl oz	PHI 1 day for green forage and silage and 28 days for grain or stover.
3, 28 R	Besiege	lambda-cyhalothrin, chlorantraniliprole	5-10 fl oz	PHI 21 days. Do not feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment. For control before the larva bores into the plant stalk or ear.
3 R	Brigade 2EC, Fanfare 2EC, Bifenture EC, Discipline 2EC,	bifenthrin	2.1-6.4 fl oz	PHI 30 days. PHI 30 days. Do not graze livestock in treated area or cut treated crops for feed within 30 days of treatment.

	Sniper, Tundra EC			
1B, 3 R	Cobalt	chlorpyrifos and gamma cyhalothrin	13-26 fl oz	PHI 21 days for grain, ears, forage or fodder.
1B, 3 R	Cobalt Advanced	chlorpyrifos and gamma cyhalothrin	11-26 fl oz	PHI 21 days for grain, ears, forage or fodder.
28	Coragen	rynaxypry	3.5-7.5 fl oz	PHI 14 days.
3 R	Declare	gamma-cyhalothrin	0.77-1.28 fl oz	PHI 21 days for grain, livestock grazing or silage harvest.
3 R	Ethos XB	bifenthrin	2.8-8.5 fl oz	PHI 30 days for grain, silage and grazing.
3 R	Fastac EC	alpha-cypermethrin	1.8-3.8 fl oz	For control before the larva bores into the plant stalk or ear. PHI 30 days for grain and stover. PHI 60 days for forage.
3 R	Hero	zeta-cypermethrin and bifenthrin	2.6-6.1 fl oz	PHI 30 days for grain or stover and 60 days for forage. Do not graze livestock in treated areas or cut treated crops for feed within 30 days of the last application.
18A	Intrepid 2F	methoxyfenozide	4-16 fl oz	PHI 21 days.
3 R	Lamcap II	lambda-cyhalothrin	0.96-1.60 fl oz	For control before the larvae bores into the ear. PHI 21 days for harvest, grazing or cutting treated crops for feed.
1B R	Lorsban Advanced, Lorsban 4E, Govern 4E, Hatchet, Nufos 4E, Warhawk, Yuma 4E	chlorpyrifos	1-2 pts	PHI 21 days before harvest of grain, ears, forage or fodder.
3 R	Mustang Maxx, Respect	zeta-cypermethrin	1.76-4.0 fl oz	Apply before the larvae enter the ear. PHI 30 days for grain and stover. PHI 60 days for forage.
1B R	PennCap-M	microencapsulated methyl parathion	2-4 pts	PHI 12 days for harvest, forage and grazing.
3 R	Pounce 25WP	permethrin	3.2-6.4 oz	PHI 30 days for grain or fodder.
28	Prevathon	chlorantraniliprole	14-20 fl oz	PHI 14 days for harvest of ears and 1 day for forage, fodder, silage or stover.
3 R	Proaxis	gamma-cyhalothrin	1.92-3.20 fl oz	For control before larvae enter the ear. PHI 21 days.
5	Radiant SC	spinetoram	3-6 fl oz	Apply broadcast or direct spray to ear zone. PHI 28 days for grain and 3 days for fodder or forage.
1A	Sevin XLR Plus	carbaryl	2 qts	PHI 48 days for grain or fodder and 14 for forage or silage.
3 R	Silencer	lambda-cyhalothrin	1.92-3.20 fl oz	PHI 21 days.

1B, 3 R	Stallion	chlorpyrifos and zeta-cypermethrin	5-11.75 fl oz	For control before the larva bores into the ear. PHI 30 days for grain or stover and 60 days for forage.
3 R	Steed	zeta-cypermethrin and bifenthrin	2.5-3.5 fl oz	PHI 30 days for grain or stover and 60 days for forage. Do not graze livestock in treated areas or cut treated crops for feed within 30 days of the last application.
22A	Steward EC	indoxacarb	6.0-11.3 fl oz	PHI 14 days for grain or stover and 1 day for forage, fodder and silage.
5 R	Tracer	spinosad	2-3 fl oz	PHI 28 days for grain and 3 days for fodder or forage harvest.
1B, 3 R	Tundra Supreme	chlorpyrifos and bifenthrin	5.6-16.8 fl oz	PHI 30 days for grain and silage.
3 R	Warrior w/ Zeon Technology	lambda-cyhalothrin	1.92-3.20 fl oz	For control before the larvae bores into the ear. PHI 21 days for harvest, grazing or cutting treated crops for feed.
3 R	Warrior II w/ Zeon Technology	lambda-cyhalothrin	1.28-1.92 fl oz	For control before the larvae bores into the ear. PHI 21 days for harvest, grazing or cutting treated crops for feed.